

# Floyd Medical Center

## Cancer Program Practice Profile Reports (CP3R)

Site	Measure Description	Measure	CoC Std/%	Estimated Performance Rates	95% CI	Compliance
Bladder	At least 2 lymph nodes are removed in patients under 80 undergoing partial or radical cystectomy (Surveillance)	BL2RLN	N/A	No data	No data	N/A
	Radical or partial cystectomy; or tri-modality therapy (local tumor destruction/excision with chemotherapy and radiation therapy) for clinical T2-4 N0 M0 patients with urothelial carcinoma of the bladder, first treatment within 90 days of diagnosis (Surveillance)	BLCSTRI	N/A	No data	No data	N/A
	Neoadjuvant or adjuvant chemotherapy recommended or administered for patients with muscle invasive cancer undergoing radical cystectomy (Surveillance)	BLCT	N/A	No data	No data	N/A
Breast	Radiation is administered within 1 year (365 days) of diagnosis for women under age 70 receiving breast conservation surgery for breast cancer (Accountability)	BCSRT	4.4/90%	94.9%	88%-100%	Compliant
	Tamoxifen or third generation aromatase inhibitor is recommended or administered within 1 year (365 days) of diagnosis for women with AJCC T1c or stage IB-III hormone receptor positive breast cancer (Accountability)	HT	4.4/90%	97.5%	92.7%-100%	Compliant
	Radiation therapy is recommended or administered following any mastectomy within 1 year (365 days) of diagnosis for breast cancer for women with $\geq 4$ positive regional lymph nodes (Accountability)	MASTRT	4.4/90%	83.3%	53.5%-100%	Compliant
	Image or palpation-guided needle biopsy to the primary site is performed to establish diagnosis of breast cancer (Quality Improvement)	nBx	4.5/80%	95.2%	90.9%-99.3%	Compliant
	Breast Conserving surgery rate for women with AJCC clinical stage 0, I, or II breast cancer (Surveillance)	BCS	N/A	86%	77%-95%	N/A
	Combination chemotherapy is recommended or administered within 4 months (120 days) of diagnosis for women under 70 with AJCC T1c N0, or stage IB-III hormone receptor negative breast cancer (Accountability)	MAC	N/A	100%	100%-100%	N/A
	Use of brachytherapy in patients treated with primary radiation with curative intent in any stage of cervical cancer (Surveillance)	CBRRT	N/A	100%	100%-100%	N/A
Cervix	Chemotherapy administered to cervical cancer patients who received radiation for stages IB-IV cancer (group 1) or with positive pelvic nodes, positive surgical margin, and/or positive parametrium (group 2) (Surveillance)	CERCT	N/A	10.8%	100%-100%	N/A
	Radiation therapy completed within 60 days of initiation of radiation among women diagnosed with any stage of cervical cancer (Surveillance)	CERRT	N/A	No data	No data	N/A
	At least 12 regional lymph nodes are removed and pathologically examined for resected colon cancer (Quality Improvement)	12RLN	4.5/85%	100%	100%-100%	Compliant
Colon	Adjuvant chemotherapy is recommended, or administered within 4 months (120 days) of diagnosis for patients under the age of 80 with AJCC stage III (lymph node positive) colon cancer (Accountability)	ACT	N/A	100%	100%-100%	N/A

# Floyd Medical Center

## Cancer Program Practice Profile Reports (CP3R) Continued

Site	Measure Description	Measure	CoC Std/%	Estimated Performance Rates	95% CI	Compliance
Endometrium	Chemotherapy and/or radiation administered to patients with stage IIIC or IV endometrial cancer (Surveillance)	ENDCTRT	N/A	No data	No data	N/A
	Endoscopic, laparoscopic, or robotic performed for all endometrial cancer (excluding sarcoma and lymphoma) for all stages except stage IV (Surveillance)	ENDLRC	N/A	40%	15.2%-64.8%	N/A
Gastric	At least 15 regional lymph nodes are removed and pathologically examined for resected gastric cancer (Quality Improvement)	G15RLN	4.5/80%	No data	No data	N/A
Kidney	At least 1 regional lymph node is removed and pathologically examined for primarily resected unilateral nephroblastoma (Surveillance)	PD1RLN	N/A	No data	No data	N/A
Lung	Systemic chemotherapy is administered within 4 months to day preoperatively or day of surgery to 6 months postoperatively, or its recommended for surgically resected cases with pathologic lymph node-positive (pN1) and (pN2) NSCLC (Quality Improvement)	LCT	4.5/85%	0%	0%-0%	Compliant
	Surgery is not the first course of treatment for cN2, M0 lung cases (Quality Improvement)	LN0Surg	4.5/85%	100%	100%-100%	Compliant
	At least 10 regional lymph nodes are removed and pathologically examined for AJCC stage IA, IB, IIA, and IIB resected NSCLC (Surveillance)	10RLN	N/A	42.9%	15.2%-64.8%	N/A
Ovary	Salpingo-oophorectomy with omentectomy, debulking/cytoreductive surgery, or pelvic exenteration in stages I-IIIC ovarian cancer (Surveillance)	OVSAL	N/A	0%	0%-0%	N/A
Rectum	Preoperative chemo and radiation are administered for clinical AJCC T3N0, T4N0, or stage III; or postoperative chemo and radiation are administered within 180 days of diagnosis for clinical AJCC T1-2N0 with pathologic AJCC T3N0, T4N0, or stage III; or treatment is recommended; for patients under the age of 80 receiving resection for rectal cancer (Quality Improvement)	RECRCT	4.5/85%	100%	100%-100%	Compliant